ES-3574 REV. 1-78



E. I. DU PONT DE NEMOURS & COMPANY

WILMINGTON, DELAWARE 19898
CENTRAL RESEARCH & DEVELOPMENT DEPARTMENT

HASKELL LABORATORY
FOR
TOXICOLOGY AND INDUSTRIAL MEDICINE

bcc: C. F. Reinhardt/B. C. McKusick

J. W. Raines - PPD - M-5625 S. G. Hundley/R. W. Hartgrove

R. E. Staples

H. J. Trochimowicz/J. G. Aftosmis

C-4124

AR 226 - 1397

March 16, 1982

Dr. Joseph Seifter
TS-792
Office of Toxic Substances
U. S. Environmental Protection Agency
401 "M" Street, S.W.
Washington, D.C. 20460

Dear Dr. Seifter:

We have studied the placental transfer of 14C-perfluoro-octanoate (C-8) in the albino rat by orally administering the chemical on Gestation Day 19 and following maternal blood and fetal tissue levels of the radiocarbon at 2, 4, and 8 hours after dosing. Maternal blood and placental levels of 14C increased between 2 and 4 hours then decreased between 4 and 8 hours after dosing. The μg equivalents in maternal blood were approximately 12, 20 and 12 $\mu g/ml$ at 2, 4, and 8 hours postosing, respectively. Corresponding fetal levels (whole body assays) were 0.7, 3, and 3 $\mu g/ml$. These data demonstrate that 14C-labelled C-8 does transfer across the placenta of the rat.

Please call me (302-366-5259) if you have any further questions.

Sincerely,

Gerald L. Kennedy, Jr. Section Supervisor, Acute Investigations

GLK: scg

HED

EID071704